

**Note to readers with disabilities:** *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact [ehp508@niehs.nih.gov](mailto:ehp508@niehs.nih.gov). Our staff will work with you to assess and meet your accessibility needs within 3 working days.

## **Supplemental Material**

### **Trihalomethanes in Drinking Water and Bladder Cancer Burden in the European Union**

Iro Evlampidou, Laia Font-Ribera, David Rojas-Rueda, Esther Gracia-Lavedan, Nathalie Costet, Neil Pearce, Paolo Vineis, Jouni J.K. Jaakkola, Francis Delloye, Konstantinos C. Makris, Euripides G. Stephanou, Sophia Kargaki, Frantisek Kozisek, Torben Sigsgaard, Birgitte Hansen, Jörg Schullehner, Ramon Nahkur, Catherine Galey, Christian Zwiener, Marta Vargha, Elena Righi, Gabriella Aggazzotti, Gunda Kalnina, Regina Grazuleviciene, Kinga Polanska, Dasa Gubkova, Katarina Bitenc, Emma H. Goslan, Manolis Kogevinas, and Cristina M. Villanueva

### **Table of Contents**

**Table S1.** Trihalomethane (THM) data collected from literature review for the countries where monitoring data were not available by national contacts.

**Table S2.** Estimated population-weighted levels of individual Trihalomethanes ( $\mu\text{g/L}$ ) in 14 European countries, 2005-2018.

**Figure S1.** Exposure-response relationship between average trihalomethane (THM) levels ( $\mu\text{g/L}$ , x-axis) and bladder cancer odds ratio (OR) (y-axis). Smoothed spline from a generalised additive model adjusted for study center, age, sex, educational level, smoking status, high-risk occupation, daily fluid intake, and coffee consumption. Tick marks above the x-axis represent observations, and the dashed lines represent the 95% confidence intervals (CIs).

**Table S3.** Estimated population attributable fraction (PAF) and number of bladder cancer (BC) cases attributable to trihalomethanes (THM) levels in 28 EU countries, men and women, age 30-79 years.

**Table S4.** Estimated number of bladder cancer (BC) cases attributable to trihalomethane (THM) levels in the lowest and highest exposure scenarios, men and women, age 30-79 years.

**Table S5.** Estimated number of attributable bladder cancer (BC) cases if no country would exceed the current EU Total Trihalomethanes (THM) mean level ( $11.7 \mu\text{g/L}$ ), men and women, age 30-79 years, EU-28.

### **References**