

Article

Efficacy of Commercial Biocontrol Products for the Management of Verticillium and Fusarium Wilt in Greenhouse Tomatoes: Impact on Disease Severity, Fruit Yield, and Quality

Maria-Dimitra Tsolakidou¹, Georgia Demetriou¹, Sotiria Panagiotou¹, Loukia Vassiliou² , Vlasios Goulas¹ 
and Iakovos Pantelides^{1,*} 

¹ Department of Agricultural Sciences, Biotechnology and Food Science, Cyprus University of Technology, Lemesos 3603, Cyprus; maria.tsolakidou@cut.ac.cy (M.-D.T.); vlasios.goulas@cut.ac.cy (V.G.)

² Department of Natural Resources and Environment, Agricultural Research Institute, Ministry of Agriculture, Nicosia 1516, Cyprus; lvassiliou@ari.moa.gov.cy

* Correspondence: iakovos.pantelides@cut.ac.cy

Abstract: *Verticillium dahliae* (Vd) and *Fusarium oxysporum* f. sp. *lycopersici* (Fol) are two major fungal pathogens that infect tomato plants, causing significant challenges in their control since both pathogens can persist in the soil for several years even in the absence of a host plant and no effective fungicides are available at present. This study investigated the efficacy of two biocontrol formulations, Clonotri (containing *Trichoderma* and *Clonostachys* microorganisms) and Strepse (comprising *Streptomyces* and *Pseudomonas* microorganisms), against Vd and Fol and their impact on tomato fruit quality and yield under greenhouse conditions. The pathogenicity experiment demonstrated that the Clonotri formulation, containing *Trichoderma* and *Clonostachys* spores, significantly reduced Fusarium wilt disease by 32% compared to the control group. However, in the Vd pathogenicity experiment, the formulations did not exhibit disease reduction, although treatment with Strepse, containing *Streptomyces* and *Pseudomonas* microorganisms, resulted in a preserved total fruit number when compared to uninfected plants. Analysis of fruit quality attributes revealed no significant differences among the various interventions. Furthermore, Fol infection in the first fruit set significantly increased fruit firmness, while Vd infection resulted in elevated levels of total soluble solids in fruits. These findings demonstrate that the evaluated biocontrol formulations provide a degree of protection against fungal wilt pathogens in tomato plants and can increase yield in greenhouse conditions while having minimal impact on overall fruit quality attributes.

Keywords: tomato; *Verticillium dahliae*; *Fusarium oxysporum* f. sp. *lycopersici*; *Trichoderma* sp.; *Clonostachys* sp.; *Streptomyces* sp.; *Pseudomonas* sp.



Citation: Tsolakidou, M.-D.; Demetriou, G.; Panagiotou, S.; Vassiliou, L.; Goulas, V.; Pantelides, I. Efficacy of Commercial Biocontrol Products for the Management of Verticillium and Fusarium Wilt in Greenhouse Tomatoes: Impact on Disease Severity, Fruit Yield, and Quality. *Agriculture* **2024**, *14*, 882. <https://doi.org/10.3390/agriculture14060882>

Academic Editor: Alessandro Vitale

Received: 8 May 2024

Revised: 28 May 2024

Accepted: 30 May 2024

Published: 1 June 2024



Copyright: © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Verticillium dahliae (Vd) and *Fusarium oxysporum* (Fox) are two of the most destructive soilborne fungal pathogens that cause wilt diseases in many economically important crops, leading to serious economic losses in production, including tomato [1]. Both pathogens infect plants by penetrating the roots, entering the xylem, and producing conidia that are transported to other parts of the plant through the xylem [2,3]. The invasion of these pathogens into the xylem vessels results in the plugging of the vessels by fungal and degraded cell wall material, disrupting water transport and causing symptoms that include wilting, yellowing of leaves, stunting, and eventually plant death [4–6].

Controlling fungal wilts caused by these pathogens is difficult due to several factors. Firstly, once plants are infected, no efficient treatments exist to prevent the progression of the disease or cure the infected plants [5,7]. Secondly, both Vd and Fox can form long-lasting survival structures such as microsclerotia and chlamydospores, respectively, which allow

them to persist in the soil for several years even in the absence of a susceptible host [8]. Resting structures represent favorable targets for control through soil solarization and chemical fumigation. Nevertheless, limitations in large-scale implementation and the prohibition of chemical fumigants due to concerns surrounding public health and environmental implications render these approaches unsuitable [9]. In addition, the application of pesticides frequently proves inadequate due to limited accessibility within the soil matrix [10]. Several strategies have been developed and implemented for the management of fungal wilts. Crop rotation, soil solarization, the use of resistant cultivars, and grafting on resistant rootstocks are commonly employed practices to reduce the incidence of *Vd* and *Fox* wilts [11–14]. Another promising strategy for managing fungal wilt pathogens is the utilization of biological control agents (BCA). BCA includes various microorganisms, such as certain strains of bacteria or fungi, that can be used to suppress the growth and activity of the pathogens through various mechanisms, including competition for nutrients and space, production of antifungal compounds, induction of systemic resistance in plants, and enhancement of plant growth and vigor [15]. When applied to the rhizosphere or as seed treatments, BCA can establish a protective presence, preventing or reducing the colonization and infection of plant roots by fungal pathogens. Additionally, the use of biological control agents offers several advantages over fungicides, including their environmental friendliness, reduced negative impacts on non-target organisms, and the potential for long-term sustainable disease management [16]. However, the effectiveness of biological control agents can vary depending on factors such as environmental conditions, application methods, and compatibility with other management practices [17]. The efficacy of biocontrol can be influenced by numerous interactions, such as variability between plants, fields, and years, and often limited efficacy and inconsistent field performance have been reported [18]. Therefore, further research is essential to optimize the use of these biological control agents under different environments and production conditions and integrate them into comprehensive disease management strategies for sustainable agriculture.

This study investigates the effectiveness of two commercially available biocontrol formulations, Clonotri and Strepse (MS BIOTECH), in managing fungal wilt pathogens, *Vd* and *Fol*, in tomato plants grown under greenhouse conditions. The primary objective is to evaluate the potential of these formulations to mitigate the disease severity caused by these fungal pathogens. Additionally, we aim to assess the impact of the biocontrol treatments on fruit yield and quality.

2. Materials and Methods

2.1. Plant Material and Growth Conditions

Tomato (*Solanum lycopersicum*) cv. Ailsa Craig (Thompson & Morgan Ltd., Ipswich, UK) was used in the experiments. Seeds were sown directly into 10.5 cm-diameter pots, each containing approximately 500 cm³ sterilized peat-based potting substrate (Plantobalt substrate 2, Plantaflor) [7]. The pots were placed in a controlled growth room (25 °C; 16 h light, 8 h dark; 65–70% RH; light intensity 300 μmol m⁻² s⁻¹ at pot height) and received watering individually every second day with an equal volume of water.

2.2. Biocontrol Formulations

In the present study, two biological formulations and their combination were evaluated against the fungal wilt pathogens of tomato. The commercial products Clonotri and Strepse (MS BIOTECH) are liquid formulations containing spores of *Trichoderma* sp., *Clonostachys* sp., and *Glomus* sp. and *Streptomyces* sp., *Pseudomonas* sp., and *Glomus* sp., respectively. According to the product labels, the formulations require activation with Nutryaction, an organic fertilizer containing yeast and brown algae extracts (MS BIOTECH), for any effect and cannot be applied directly for plant protection.

The following treatment solutions were prepared:

Single Application of Clonotri: This solution was prepared by mixing 1 part Clonotri formulation with 13 parts water and 2 parts Nutryaction (*v/v*) at least 8 h prior to application on plants.

Single Application of Strepse: This solution was prepared following the same protocol as the single application of Clonotri, using 1 part Strepse formulation instead.

Combined Application of Clonotri and Strepse: To evaluate the potential synergistic effects, a combined application was prepared using half the recommended dosage of each individual product. This solution was prepared by mixing 0.5 part Clonotri formulation, 0.5 part Strepse formulation, 13 parts water, and 2 parts Nutryaction (*v/v*) at least 8 h before application.

2.3. Fungal Strains and Inoculum Preparation

The fungal isolates of *Vd* (race 1 isolate 70 V) [19] and *Fol* (race 1 isolate Fol004) [20] used in this study were obtained from the culture collection maintained at the Department of Agricultural Sciences, Biotechnology, and Food Science, Cyprus University of Technology, Limassol, Cyprus. Fungal isolates were cryopreserved in an aqueous 20% glycerol suspension at $-80\text{ }^{\circ}\text{C}$. Before being used, fungal strains were activated on potato dextrose agar (PDA, Merck, Rahway, NJ, USA) at $25\text{ }^{\circ}\text{C}$ for 5 days. For the pathogenicity experiments, conidia were prepared by transferring pieces from the growing edge of the fungal colony of each pathogen in sucrose sodium nitrate (SSN) [21] to 500 mL Erlenmeyer flasks for the growth room experiments and 2-L Erlenmeyer flasks for the greenhouse experiments. The flasks were then incubated in an orbital shaker at 150 rpm in the dark for 5 days. *Vd* was grown at $22\text{ }^{\circ}\text{C}$, and *Fol* at $25\text{ }^{\circ}\text{C}$. Liquid cultures were passed through cheesecloth to remove mycelia. Suspensions were centrifuged at $10,000\times g$ at $8\text{ }^{\circ}\text{C}$ for 10 min and resuspended in sterile distilled water (SDW).

2.4. Pathogenicity Experiments in Growth Room

We performed two separate growth room experiments. In the first pathogenicity experiment, the two formulations were initially evaluated under growth room conditions against *Vd* and *Fol*. The treatments for this experiment were as follows:

(a) Negative Control (-ve Control): Plants grown in peat-based potting substrate only, serve as the healthy control group. Peat is a standard choice in plant growth studies investigating disease control as it reflects a scenario where plants are not treated with any additives or biocontrol products.

(b) Strepse: Plants grown in peat-based potting substrate (Peat) and treated with activated Strepse formulation.

(c) Clonotri: Plants grown in Peat and treated with activated Clonotri formulation.

(d) Mix: Plants grown in Peat and treated with a mixture of activated Strepse and Clonotri formulation.

(e) *Vd*+ control: Plants grown in Peat and inoculated with *Vd*, serving as the infected, untreated control group. This group allows for comparison of *Vd* disease severity in the absence of any protective measures.

(f) Strepse *Vd*: Plants grown in Peat, treated with activated Strepse formulation and inoculated with *Vd*.

(g) Clonotri *Vd*: Plants grown in Peat, treated with activated Clonotri formulation, and inoculated with *Vd*.

(h) Mix *Vd*: Plants grown in Peat, treated with activated Strepse and Clonotri formulation, and inoculated with *Vd*.

(i) *Fol*+ control: Plants grown in Peat and inoculated with *Fol*, serving as the infected, untreated control group. This group allows for comparison of *Fol* disease severity in the absence of any protective measures.

(j) Strepse *Fol*: Plants grown in Peat, treated with activated Strepse formulation, and inoculated with *Fol*.

(k) Clonotri *Fol*: Plants grown in Peat, treated with activated Clonotri formulation, and inoculated with *Fol*.

(l) Mix *Fol*: Plants grown in Peat, treated with activated Strepse and Clonotri formulation, and inoculated with *Fol*.

Application of treatments was performed twice via root drenching, 7 and 14 days after seedling emergence (10 mL/plant in each application). Control plants were root-drenched with 10 mL of H₂O only. Plants were artificially inoculated at the third leaf stage (~20 days after seedling emergence) by root drenching with 20 mL of conidial suspension of *Vd* or *Fol*. This method involved pouring the conidial suspension directly onto the root zone of the plants in their pots. Conidial suspension concentration was determined using a hemocytometer, allowing for accurate quantification of spores before applying the inoculum to the plants. The suspensions were diluted appropriately in sterile distilled water to achieve the desired final concentrations of 10⁶ conidia/mL for *Vd* and 10⁷ conidia/mL for *Fol*. Control plants were mock-inoculated with 20 mL of sterile distilled water. Plants were arranged in rows and columns in a completely randomized design and were rotated within the growth room every second day before watering. The plants were watered individually every second day with an equal volume of water.

Disease severity was calculated by the number of leaves showing typical symptoms (wilting, yellowing, and browning of leaves) as a percentage of the total number of leaves of each plant. Symptoms were periodically recorded for 19 days post-inoculation (dpi) for both pathogens. Disease ratings were plotted over time to generate disease progress curves. The area under the disease progress curve (AUDPC) was calculated by the trapezoidal integration method [22]. Disease was expressed as a percentage of the maximum possible area for the whole period of the experiment, referred to as relative AUDPC. The experiment was repeated three times, with 10 replicates per treatment.

In the second experiment, based on the results of the first pathogenicity experiment, Strepse was evaluated against *Vd*, and Clonotri was evaluated against *Fol*. In this experiment, the efficacy of the products was assessed at three different pathogen concentrations (10⁶, 10⁵, and 10⁴ conidia/mL), and adjusted using a hemocytometer to reach the desired levels. The remaining experimental procedure was performed as described above. Apart from disease severity, plant height and total biomass of the plants were determined at 22 dpi (the last day of disease symptom monitoring).

2.5. Pathogenicity Experiments in Greenhouse

The efficacy of the biocontrol formulations was also evaluated under greenhouse conditions against *Vd* and *Fol* with pathogenicity experiments on tomato plants. This experiment was carried out in a greenhouse at the Agricultural Research Institute (35.140416, 33.383872, Nicosia, Cyprus) during the period from 1 November 2019, to 10 March 2020. The treatments of this experiment were the same as the first pathogenicity experiment in the growth room (described in Section 2.4), with 8 replicates per treatment. This experiment represents data collected during a single growing season (November 2019–March 2020). Tomato plant cv. Ailsa Craig was initially grown in a controlled growth room as described previously (Section 2.1) and received the biocontrol treatments via root drenching, 7 and 14 days after seedling emergence (10 mL/plant in each application). Control plants were root-drenched with 10 mL of H₂O only. Plants were transferred to the greenhouse and were artificially inoculated with *Vd* or *Fol* at the third leaf stage (~20 days after seedling emergence) by root dipping. Briefly, the plants were removed carefully from their original pots while preserving the root ball. They were then immersed in a tray containing conidial suspension of *Vd* or *Fol*, adjusted to 10⁶ conidia/mL and 10⁷ conidia/mL, respectively. Each plant remained in the tray for 10 min. This immersion method ensured a more thorough exposure of the entire root system to the pathogen inoculum. Control plants were immersed in H₂O for 10 min. Then the plants were transplanted into 15 L pots with a 30 cm diameter and filled with 3 kg of peat-based potting substrate (Plantobalt substrate 2, Plantaflor). The pots were arranged in rows and columns in a completely randomized

design, and the plants were monitored for disease development and symptoms caused by *Vd* or *Fol* infection. Disease severity was assessed based on the percentage of leaves showing typical symptoms, and the Area Under the Disease Progress Curve (AUDPC) was calculated using the trapezoidal integration method as described in detail in Section 2.4 [22]. Symptoms were periodically recorded for 52- and 50-days post-inoculation (dpi) for *Vd* and *Fol*, respectively.

The temperature inside the greenhouse during the experiment ranged between 14 °C and 27 °C. The temperature control within this range was achieved through appropriate greenhouse control systems (cooling/heating systems, humidity sensors, ventilation fans). In the initial stages of cultivation, the plants were irrigated daily, and as time progressed, they were irrigated every two days according to their water needs (the volume of irrigation water ranged from 600 mL during the establishment stage to 1 L during the harvesting stage). This approach prevented excessive water drainage from the pots. Throughout the experiment, the plants were subjected to two additional interventions with the biocontrol formulations at 20-day intervals after pathogen inoculation by mixing the biocontrol products with the irrigation water. During plant growth, vertical support was provided to the plants using greenhouse trellises, and any lateral shoots were promptly removed upon emergence. The onset of the first flowering was observed around 35 days after transplanting the plants in the greenhouse, followed by the appearance of the initial tomato fruits approximately one to two weeks later (around 56 days after transplantation). Fertilization of the plants was applied twice during the entire experiment using Betabio Full organic fertilizer (ED&F Man Liquid Products MLP) mixed with the irrigation water, following the recommendations of the company that provided the biocontrol formulations for this specific greenhouse setup.

2.6. Quantity Attributes

A total number of fruits per plant was recorded for each treatment. The fruit was harvested, and the total fresh weight of the fruit was determined for each plant using a digital scale with an approximation of 0.01 g. Plant height was measured for each plant on the last day of the experiment (131 days after transplant) as the height above ground where the youngest fully expanded leaf joined the stem. The fresh weight of plants was measured for each plant 131 days after transplant using a digital scale with an approximation of 0.01 g. The number of leaves was counted for each plant 131 days after transplant (including removed, dead, or fallen leaves).

2.7. Determination of Quality Attributes

The quality attributes of tomato fruits were determined in tomato fruits as described in previous work [23]. Tomato fruits were harvested from the first two fruit sets, 117 and 131 days after transplantation, in 15 L pots.

The external fruit color was assessed at two equatorial areas of each fruit using a Konica Minolta CR-400 chromameter (Konica Minolta Sensing, Inc., Osaka, Japan). The equipment was set up for a D65 illuminant and a 10° observer angle. A white standard plate ($Y = 94.00$, $x = 0.3158$, $y = 0.3322$) was used for calibration. The values of L^* (lightness), a^* (green to red color), and b^* (blue to yellow color) were determined for each treatment.

Firmness was evaluated by measuring the maximum penetration force for a 3-mm-diameter steel probe using a TA-XT2 texture analyzer (Stable Micro Systems Ltd., Surrey, UK). The fruits were placed so that the plunger penetrated the pericarp approximately 1 cm away from their geometric center to a depth of 10 mm at a rate of 5 mm/s. Results were expressed in Newtons (N).

Tomato fruits from each treatment were homogenized in a blender and centrifuged in order to obtain a clear juice. Soluble solids content (SSC) was measured with a portable digital refractometer, the ATAGO Palette PR-32a (Atago Company Ltd., Tokyo, Japan). Titratable acidity (TA) was determined by potentiometric titration with 0.1 mol/L NaOH up to pH 8.2, using 5 mL of juice diluted in distilled water until final volume of 50 mL. The

measurements were carried out using a DL22 Mettler Toledo titrator (Mettler–Toledo, Inc., Columbus, OH, USA) and were expressed as g citric acid L⁻¹.

2.8. Determination of Total Phenolic and Total Flavonoid Contents

The phenolic contents of tomato fruits were extracted using ultrasound-assisted extraction according to Ochoa-Velasco et al. [24]. Approximately 5 g of tomato pulp was mixed with 20 mL methanol/water (80:20, *v/v*). The mixture was placed into an ultrasound bath for 30 min at 30 °C. After extraction, the mixture was centrifuged and the supernatant was collected to determine phenolic and flavonoid contents as well as the antioxidant activity [24].

The total phenolic contents were estimated using a microplate Folin–Ciocalteu assay. More specifically, 50 µL of extracts were mixed with 50 µL of Folin–Ciocalteu reagent (1:5, *v/v*) and 100 µL of sodium hydroxide solution (0.35 mol/L) in each well. After a period of 3 min, the absorbance of each sample was monitored at 760 nm. A standard curve of gallic acid was prepared and results were expressed as mg gallic acid equivalents (GAE)/100 g tomato fruit.

The total flavonoid contents were also determined with the use of a microplate-adopted photometric assay. An aliquot of 25 µL of the sample or standard solution was mixed with 100 µL distilled water and 10 µL of 50 g/L NaNO₂. After 5 min, 15 µL of 100 g/L AlCl₃ was added to the mixture. Finally, aliquots of 50 µL of sodium hydroxide solution (1 mol/L) and 50 µL of distilled water were added after 6 min. The mixture was shaken for 30 s in the plate reader prior to absorbance measurements at 510 nm. A standard curve of rutin hydrate was prepared, and the results were expressed as mg rutin equivalents (RE)/100 g tomato fruit [25].

2.9. Determination of β-Carotene and Lycopene Contents

Sixteen mL of acetone-hexane (4:6) solvent was added to 1 g of tomato pulp and thoroughly mixed with the use of a homogenizer (Ultra–Turrax, IKA T25, Staufen, Germany). When the two phases were separated, an aliquot was taken from the upper phase, and the absorbance at 663, 645, 505, and 453 nm was determined. Beta-carotene and lycopene contents were calculated according to the Nagata and Yamashita equations and expressed as mg/L [26].

2.10. Determination of Antioxidant Activity

The 1,1-diphenyl-2-picrylhydrazyl (DPPH) assay was used to measure the scavenging activity of antioxidants against the colorful radicals of DPPH. The reaction mixtures consisted of 30 µL of extracts and 200 µL DPPH methanolic solution (0.2 mM). The mixtures were incubated for 30 min; then, the absorbance was read at 515 nm. A standard curve of 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (Trolox) was prepared, and results expressed as µmol Trolox equivalent/100 g tomato fruit.

The antioxidant activity of the polar fraction of tomato fruits was also tested using Ferric Reducing Antioxidant Power (FRAP) measurements. More specifically, 100 µL of the sample or standard compound were mixed with 180 µL FRAP solution (300 mM acetate buffer at pH 3.6, 10 mM 2,4,6-Tris(2-pyridyl)-s-triazine solution, and 20 mM ferric chloride solution in the ratio of 10:1:1, *v/v/v*) in 96-well plates. The reaction mixtures in well plates were allowed for 6 min at 37 °C and the absorbance was read at 595 nm. A standard curve of iron sulfate (FeSO₄) was prepared, and the results were expressed as µmol FeSO₄/100 g of tomato fruit [24].

2.11. Statistical Analysis

Statistical analyses were performed using GraphPad Prism 8 for Windows (GraphPad Software, La Jolla, CA, USA). The type of ANOVA used depended on the experimental design.

A one-way ANOVA was used for pathogenicity experiments where data were collected from groups with a single factor. Following the ANOVA, Tukey's post-hoc test ($p \leq 0.05$) was used for multiple comparisons between groups.

A two-way ANOVA was employed for experiments where the data were influenced by two independent factors. In these cases, the interaction effect between the two factors was also assessed. If the interaction was significant, Tukey's post-hoc test with adjustment for multiple comparisons ($p \leq 0.05$) was used to identify specific significant differences between groups.

3. Results

3.1. Evaluation of Biocontrol Products against Fungal wilt Pathogens of Tomato in Controlled Conditions

In this experiment, the effect of the biocontrol formulations Clonotri (*Trichoderma* and *Clonostachys*-based product) and Strepse (*Streptomyces* and *Pseudomonas*-based product) was evaluated against *Vd* and *Fol* on tomato plants. Initial symptoms induced by *Vd* inoculation were observed on the 11th day post-inoculation (dpi). Symptoms of the disease (in the form of wilting followed by yellowing and necrosis of leaves) were recorded until 19 dpi (Figure 1a). The treatment with Strepse resulted in slower disease development and milder symptoms compared to non-treated control plants (plants that received no pathogen or biocontrol formulation application). Similarly, plants treated with Clonotri and the mixture of Clonotri and Strepse (Mix) displayed an intermediate disease severity between the Strepse treatment and the non-treated control plants (Figure 1a). The relative Area Under Disease Progress Curve (AUDPC) [22] values over a 19-day period were 40% lower in Strepse-treated plants (Strepse *Vd*) compared to the water-treated plants (*Vd*+ control). However, there were no significant differences in relative AUDPC values among the Strepse-treated (Strepse *Vd*), Clonotri-treated (Clonotri *Vd*), and Mix-treated (Mix *Vd*) plants (Figure 1b).

Tomato bioassays with *Fol* revealed initial symptoms at 11 dpi. These symptoms included wilting, chlorosis, and yellowing, and were most pronounced on older leaves. The symptoms were recorded until 19 dpi. Plants treated with Clonotri (Clonotri *Fol*) showed fewer prominent symptoms and slower disease development compared to the other treatments. At 19 dpi, the Clonotri *Fol* plants exhibited a disease severity of 21.9%, while those treated with Strepse (Strepse *Fol*), Mix (Mix *Fol*), and water (*Fol*+ control) displayed disease severities of 46.8%, 33.6%, and 39.3%, respectively (Figure 1c). The relative AUDPC values over 19 days of disease progress were higher in *Fol*+ control (15.2%), Strepse *Fol* (16.4%), and Mix *Fol* plants (14.3%) compared to Clonotri *Fol* plants (6.0%) (Figure 1d).




Figure 1. Pathogenicity experiment in controlled conditions: (a) Disease progress over time on tomato plants infected with *Verticillium dahliae*; (b) Amount of *Verticillium dahliae* disease expressed as relative AUDPC; (c) Disease progress over time on tomato plants infected with *Fusarium oxysporum* f. sp. *lycopersici*; (d) Amount of *Fusarium oxysporum* f. sp. *lycopersici* disease expressed as relative AUDPC. Vertical bars indicate the SE of the mean ($n = 10$ replicates). Different letters indicate significant differences: $p < 0.05$; one-way ANOVA, Tukey's test. Vd+ control: Plants grown in peat-based potting substrate (Peat) and inoculated with Vd. Strepse Vd: Plants grown in Peat, received activated Strepse formulation and inoculated with Vd. Clonotri Vd: Plants grown in Peat, received activated Clonotri formulation and inoculated with Vd. Mix Vd: Plants grown in Peat, received activated Strepse and Clonotri formulation and inoculated with Vd. Fol+ control: Plants grown in Peat and inoculated with Fol. Strepse Fol: Plants grown in Peat, received activated Strepse formulation and inoculated with Fol. Clonotri Fol: Plants grown in Peat, received activated Clonotri formulation and inoculated with Fol. Mix Fol: Plants grown in Peat, received activated Strepse and Clonotri formulation and inoculated with Fol.

3.2. Evaluation of Biocontrol Products against Varied Pathogen Concentrations in a Controlled Environment

In light of the results obtained in the previous experiment, the biocontrol products were evaluated against varying concentrations of Vd and Fol. Strepse was subjected to evaluation against a range of concentrations of Vd, while Clonotri underwent assessment against varying concentrations of Fol. The disease progression for both pathogens exhibited a similar pattern as observed in the previous experiment, with initial symptoms appearing at 12 dpi for Vd and 14 dpi for Fol. The symptoms were periodically recorded until 22 dpi (Figures 2a–c and 3a–c). More specifically, regarding Vd, AUDPC analysis demonstrated a significant interaction effect, indicating that the effectiveness of Strepse in reducing disease severity depended on the concentration of the pathogen. The interaction plot (Figure S1) revealed that the Strepse effect on disease severity changed across the three pathogen concentrations and was more effective at 10^5 and 10^6 conidia/mL. Post-hoc analysis comparing disease severity between Strepse-treated and control groups at each pathogen concentration level separately revealed that Strepse provided substantial protection to plants at pathogen concentrations of 10^5 and 10^6 conidia/mL, whereas the level of protection was not significant at a concentration of 10^4 conidia/mL (Figure 2d).




Figure 2. Pathogenicity experiment against varied pathogen concentrations of *Verticillium dahliae* (Vd): (a–c) Disease progress over time on tomato plants inoculated with 10^4 , 10^5 , and 10^6 conidia/mL of the pathogen, respectively; (d) Amount of disease expressed as relative AUDPC. Vertical bars indicate the SE of the mean ($n = 10$ replicates). Different letters indicate significant differences: $p < 0.05$; two-way ANOVA, Tukey’s test.




Figure 3. Pathogenicity experiment against varied pathogen concentrations of *Fusarium oxysporum* f. sp. *lycopersici* (Fol): (a–c) Disease progress over time on tomato plants inoculated with 10^4 , 10^5 , and 10^6 conidia/mL of the pathogen, respectively; (d) Amount of disease expressed as relative AUDPC. Vertical bars indicate the SE of the mean ($n = 10$ replicates). Different letters indicate significant differences: $p < 0.05$; two-way ANOVA, Tukey’s test.

Concerning *Fol*, the analysis of AUDPC showed a significant interaction, suggesting that the ability of Clonotri to reduce disease severity depended on the concentration of the pathogen. As seen in Figure S2, the impact of Clonotri on disease severity varied across the three pathogen concentrations and was more effective at 10^6 conidia/mL. Post-hoc analysis comparing disease severity between Clonotri-treated and control groups at each pathogen concentration level individually showed that Clonotri provided significant protection to plants only at a pathogen concentration of 10^6 conidia/mL, whereas the level of protection was not significant at concentrations of 10^4 and 10^5 conidia/mL (Figure 3d).

In this experiment, the height and fresh weight of plants were also measured. The application of the biocontrol products did not have any significant effect on these parameters, regardless of whether the plants were part of the control group or were inoculated with a pathogen (Figures S3 and S4).

3.3. Evaluation of Biocontrol Products in Greenhouse Conditions

In an attempt to assess the biocontrol products under more practical conditions, a pathogenicity experiment was carried out in a greenhouse. In addition to evaluating their efficacy against disease development, the biocontrol products were also analyzed for their impact on fruit yield and quality.

3.3.1. Pathogenicity Experiment

Strepse and Clonotri were subsequently assessed for their effectiveness against *Vd* and *Fol* in the greenhouse experiment. This evaluation aimed to determine whether their efficacy remained consistent when challenged by these specific pathogens under different environmental conditions. To ensure a more pronounced and measurable impact of the biocontrol products on disease reduction, a concentration of 10^6 conidia/mL was selected for both pathogens in the greenhouse experiment. This concentration allowed for a clearer evaluation of the biocontrol products' effectiveness by providing a stronger pathogenic challenge that could be more effectively countered by the biocontrol agents.

In the *Vd* bioassay, the first symptoms were observed at 6 dpi in the form of wilting and chlorosis, particularly on older leaves, and symptoms were periodically recorded until 52 dpi (Figure 4a). Although the Strepse *Vd* plants exhibited fewer symptoms at nearly all recorded time points compared to the other interventions, AUDPC analysis revealed that this difference was not significant (Figure 4b).




Figure 4. Cont.





Figure 4. Pathogenicity experiment in greenhouse conditions: (a) Disease progress over time on tomato plants infected with *Verticillium dahliae* (*Vd*); (b) Amount of *Vd* disease expressed as relative AUDPC; (c) Disease progress over time on tomato plants infected with *Fusarium oxysporum* f. sp. *lycopersici* (*Fol*); (d) Amount of *Fol* disease expressed as relative AUDPC. Vertical bars indicate the SE of the mean ($n = 8$ replicates). Different letters indicate significant differences: $p < 0.05$; one-way ANOVA, Tukey’s test. (e) Representative photographs of tomato plants at 50 days post-inoculation with *Vd* (up) and *Fol* (down). Plants inoculated with *Vd* exhibit similar wilting symptoms across all treatments, whereas the Clonotri *Fol* treatment shows significantly reduced wilting compared to the other *Fol* treatments.

In the pathogenicity experiment with *Fol*, initial symptoms of wilting and chlorosis were observed at 11 dpi. Symptom development was then monitored periodically until 50 dpi (Figure 4c). Disease symptoms progressed similarly in the *Fol*+ control, Strepse *Fol*, and Mix *Fol* plants. However, Clonotri *Fol* plants exhibited fewer symptoms and reduced disease severity at most time points (Figure 4c). For instance, at 22 dpi, Clonotri *Fol* plants showed a disease incidence of 60%, while the other treatments displayed a disease incidence of approximately 95–100%. Clonotri *Fol* treatment significantly reduced AUDPC by 32% compared to the *Fol*+ control and by 33% and 37% compared to Strepse *Fol* and Mix *Fol*, respectively. However, no significant difference was observed among the latter three treatments ($p < 0.05$) (Figure 4d).

3.3.2. Plant Growth Traits

In the greenhouse experiment, the effects of Strepse and Clonotri were assessed for plant growth characteristics. Plant height, number of leaves, and plant fresh weight were measured in both control plants and plants infected with *Vd* and *Fol*. A two-way ANOVA was performed to analyze the effect of biocontrol application and pathogen inoculation on these parameters. The analysis revealed that there was not a statistically significant interaction between the effects of biocontrol application and pathogen inoculation on plant fresh weight ($F = 1.39$, $p = 0.23$), plant height ($F = 0.78$, $p = 0.59$), and number of leaves ($F = 1.16$, $p = 0.34$). Simple main effects analysis showed that biocontrol application did not have a statistically significant effect on plant fresh weight ($p = 0.23$), plant height ($p = 0.59$), or number of leaves ($p = 0.34$), whereas pathogen inoculation had a statistically significant effect on plant weight ($p = 0.001$) but not on plant height ($p = 0.28$) or number of leaves ($p = 0.14$). As shown in Figure S5, plant fresh weight was significantly higher in control plants treated with Strepse (Strepse-treated control plants) compared to plants inoculated with the pathogens.

In terms of production characteristics, the total number of fruits, mean fruit fresh weight, and overall productivity (measured as the total fresh weight of harvested tomato fruits) were evaluated in both control plants and plants infected with *Vd* and *Fol*. Two-way ANOVA analysis showed there was a significant interaction between the effects of biocontrol application and pathogen inoculation only on mean fresh weight ($F = 2.54$, $p = 0.02$) and not on number of fruits ($F = 0.64$, $p = 0.70$) or productivity ($F = 0.73$, $p = 0.63$). Simple main effects analysis showed that biocontrol application did not have a statistically significant effect on production characteristics (mean fresh weight, $p = 0.08$; fruit number, $p = 0.23$; productivity, $p = 0.35$) whereas pathogen inoculation had a significant effect on all parameters evaluated (plant weight, $p = 0.001$; plant height, $p = 0.28$; number of leaves, $p = 0.14$). Post-hoc analysis on mean fresh weight showed no significant difference between Clonotri *Fol* plants compared to the mock-inoculated plants treated with Clonotri (Clonotri-treated control plants); however, in the remaining treatments, the application of biocontrol products did not provide protection against any decline in fruit fresh weight (Figure 5b). Simple effects within pathogen inoculation showed that Strepse *Vd* treatment did not significantly impact overall fruit number compared to non-infected plants; however, none of the other treatments effectively protected against a decline in number of fruits (Figure 5a). Finally, none of the treatments successfully protected overall productivity in plants infected with the pathogens (Figure 5c).

In summary, a significant interaction between biocontrol and pathogen inoculation was only observed for mean fruit weight. Clonotri provided some protection against a decline in mean fruit weight for *Fol* infection, but overall, neither biocontrol agent significantly improved fruit production characteristics (number of fruits, mean fruit weight, and overall productivity) in plants infected with the pathogens.




Figure 5. Production characteristics in greenhouse experiment: (a) Fruits number on *F. oxysporum* f. sp. *lycopersici* (Fol) and *Verticillium dahliae* (Vd) inoculated plants; (b) Mean Fresh Weight on Fol and Vd inoculated plants; (c) Total Productivity (measured as total fresh weight of harvested tomato fruits) on Fol and Vd inoculated plants. Vertical bars indicate the SE of the mean (n = 8 replicates). Different letters indicate significant differences: p < 0.05; two-way ANOVA, Tukey's test. -ve control: plants grown in peat-based potting substrate only, serving as the control treatment. Strepse: plants grown in Peat that received activated Strepse formulation. Clonotri: plants grown in Peat that received activated Clonotri formulation. Mix: plants grown in Peat that received a mixture of activated Strepse and Clonotri formulation.

3.3.3. Fruit Quality Attributes

The results of fruit firmness were analyzed with a two-way ANOVA and showed that there was no significant interaction between the effects of biocontrol application and pathogen inoculation on the tomatoes of the first ($F = 2.09, p = 0.05$) and second fruit sets ($F = 2.40, p = 0.07$). Simple main effects analysis showed that biocontrol application did not have a statistically significant effect on fruit firmness of the first ($p = 0.17$) and the second fruit set ($p = 0.07$), indicating that the biocontrol products tested in this experiment did not have a direct impact on the fruit firmness under the conditions of the study. Conversely, simple main effects analysis showed that pathogen inoculation had a significant effect on both fruit sets ($p < 0.001$). In the *Fol*-inoculated treatments of the first fruit set, fruit firmness increased compared to the mock-inoculated plants in all treatments. Conversely, fruit firmness remained unaffected in *Vd*-inoculated plants compared to the mock-inoculated ones (Figure 6). However, this effect was not observed in the samples of the second fruit set (Figure S6).




Figure 6. Fruit firmness of the first fruit set in plants inoculated with *V. dahliae* and *F. oxysporum* f. sp. *lycopersici* and treated with biocontrol products in the greenhouse experiment. Vertical bars indicate the SE of the mean. Different letters indicate significant differences: $p < 0.05$; two-way ANOVA, Tukey's test.

External fruit color exhibited a similar trend to that observed for fruit firmness. A two-way ANOVA analysis revealed a significant interaction between the effects of biocontrol application and pathogen inoculation on the external color of the tomatoes of the first set but not of those of the second fruit set. In *Fol*-inoculated treatments, a significant increase in lightness (L^* values) was observed in the tomatoes of the first set compared to other treatments, suggesting that infection of plants with *Fol* leads to a lighter appearance of the fruit. The values for the parameters a^* (red-green color component) and b^* (yellow-blue color component) varied among the treatments. In *Fol*-inoculated plants that were treated with biocontrol products and their mixture, samples showed negative a^* values and higher positive b^* values, indicating a shift towards green and yellow coloration compared to the other treatments (Table 1). The results for external fruit color, similar to fruit firmness, did not show any significant impact of *Fol* in the samples of the second fruit set (Table S1). It is also worth mentioning that the application of the biocontrol products and their mixture did not demonstrate any significant effect on the external fruit color of the control plants (Table 1).

Table 1. Mean values of color attributes L* (lightness), a* (green to red color), and b* (blue to yellow color) in tomato fruit of the first fruit set, as affected by the biocontrol products and the pathogens.

Treatment	L* (Lightness)	a* (Green to Red Color)	b* (Blue to Yellow Color)
Negative control	41.63 ± 0.79 de	12.48 ± 1.28 bcd	20.57 ± 0.46 e
Fol+ control	47.45 ± 1.18 abc	3.56 ± 3.09 de	24.68 ± 0.70 abc
Vd+ control	43.66 ± 1.09 bcde	16.52 ± 1.98 ab	24.1 ± 0.87 abcd
Clonotri	43.24 ± 0.90 cde	10.83 ± 1.46 bcd	21.78 ± 0.43 cde
Clonotri Fol	50.69 ± 0.72 a	−6.72 ± 1.85 f	25.09 ± 0.35 ab
Clonotri Vd	43.21 ± 1.01 cde	14.61 ± 2.01 ab	22.87 ± 0.68 abcde
Strepse	40.79 ± 0.80 de	15.05 ± 1.39 ab	21.16 ± 0.47 de
Strepse Fol	51.06 ± 1.20 a	−4.60 ± 2.58 ef	24.47 ± 0.62 abc
Strepse Vd	45.27 ± 1.20 bcd	13.07 ± 2.10 bc	25.07 ± 0.88 ab
Mix	42.30 ± 0.90 de	12.00 ± 1.50 bcd	20.94 ± 0.45 e
Mix Fol	48.50 ± 1.65 ab	4.96 ± 3.45 cd	25.33 ± 0.84 a
Mix Vd	40.36 ± 0.51 e	23.17 ± 0.56 a	22.16 ± 0.63 abc

Values with different letters within each column are significantly different: $p < 0.05$; two-way ANOVA, Tukey’s test.

The results of soluble solids content (SSC) were analyzed with a two-way ANOVA and showed that there was no significant interaction between the effects of biocontrol application and pathogen inoculation on the SSC of the tomatoes of the first ($F = 0.36, p = 0.90$) and second fruit sets ($F = 1.35, p = 0.27$). Simple main effects analysis showed that biocontrol application did not have a statistically significant effect on the SSC of the first ($p = 0.43$) and the second fruit set ($p = 0.81$), indicating that the biocontrol products tested in this experiment did not have a direct impact on fruit firmness under the conditions of the study. Conversely, simple main effects analysis showed that pathogen inoculation had a significant effect on the SSC of both fruit sets ($p < 0.001$). Simple effects within biocontrol treatment showed that the soluble solids content was increased in all plants inoculated with *Vd* (Figure 7a,b) suggesting that the *Vd* infection might enhance certain metabolic processes related to fruit ripening or sugar accumulation. The soluble solids content was not affected in *Fol*-inoculated plants or plants treated with the biocontrol products (Figure 7).




Figure 7. Soluble solids content (SSC) in the fruits of tomato plants inoculated with *V. dahliae* and *F. oxysporum* f. sp. *lycopersici* and treated with biocontrol products in the greenhouse: (a) SSC of the first fruit set; (b) SSC of the second fruit set. in. Vertical bars indicate the SE of the mean. Different letters indicate significant differences: $p < 0.05$; two-way ANOVA, Tukey’s test.

No significant differences were observed in titratable acidity, total phenolic and total flavonoid contents, β -carotene and lycopene contents, or antioxidant activity (measured using DPPH and FRAP assays) among the different treatments (Figure S7). These data indicate that the application of biocontrol products and the presence of the pathogens (*Fol* and *Vd*) did not have a substantial impact on these particular biochemical parameters in the plants that are linked with organoleptic attributes and the nutritional value of fruits.

4. Discussion

Building upon the established knowledge of *Verticillium dahliae* (*Vd*) and *Fusarium oxysporum* f. sp. *lycopersici* (*Fol*) as detrimental soilborne pathogens for tomato cultivation [1], this study investigated the effectiveness of two biocontrol formulations, Strepse and Clonotri, in managing these pathogens under controlled conditions and greenhouse experiments.

Clonotri, a formulation comprising *Trichoderma* and *Clonostachys* microorganisms, exhibited significant protection against *Fol* in both controlled conditions and the greenhouse, as indicated by reduced disease severity and relative AUDPC values (Figure 1c,d and Figure 4c,d). This is consistent with previous studies that have highlighted the role of *Trichoderma* and *Clonostachys* species in suppressing *Fol*, both in controlled conditions and greenhouse environments. Through a sequence of greenhouse experiments, the introduction of non-pathogenic strains of *Trichoderma* into a potting mix resulted in substantial disease control against *Fusarium* wilt in tomato seedlings, with similarly effective outcomes observed using commercially available biocontrol products containing *Trichoderma* spp. [27]. Likewise, the effectiveness of *Trichoderma asperellum* strain T34 against *Fol* on tomato plants was demonstrated within a controlled environmental chamber experiment, revealing its efficacy across varying iron concentrations and its capacity to significantly reduce *Fol* populations while safeguarding plants from *Fusarium* wilt [28]. Another study highlighted the potential of *Trichoderma* isolates to augment the suppressive capacity of growth media in combating *Fusarium* wilt of tomato, underscoring their ability to effectively mitigate the disease during the initial phases of plant development [29]. Moreover, *Trichoderma asperellum* demonstrated robust antifungal activity against *Fol*, employing mechanisms such as chitinase activity and bioactive compound production while also stimulating plant growth and fortifying tomato plants' resistance to *Fusarium* wilt [30]. Additionally, in a growth chamber study, *Clonostachys rosea* isolate IK726 demonstrated effective control of seed- and soil-borne diseases, such as *Fusarium* wilt. This effectiveness was particularly evident when roots were pre-inoculated with *Clonostachys*. This finding highlights its capacity for systemic colonization, extending from roots to stems. [31].

Clonotri, assessed in both growth room and greenhouse experiments, did not demonstrate effectiveness against *Vd*. Despite the conducted trials, the biocontrol agent did not display significant control over the pathogen in either controlled environment (Figures 1a,b and 4a,b). The efficacy of *Trichoderma*-based treatments against *Vd* infections appears to exhibit variable outcomes, as evidenced by the findings from various studies, including our own. In some instances, *Trichoderma* interventions have demonstrated significant effectiveness in reducing *Vd* colonization and subsequently improving crop yield. A study conducted over two consecutive years in potato fields demonstrated that the integration of *Trichoderma harzianum* with other treatments led to a substantial reduction in *Vd* colonization of stems, resulting in notable increases in potato yield. This trend was consistent across different cultivars, further emphasizing the potential of *Trichoderma*-based strategies [32]. Conversely, another study exploring *Trichoderma virens* strains for their ability to induce resistance against *Verticillium* wilt in cotton plants revealed a more nuanced scenario. While certain strains of *T. virens* exhibited a significant reduction in disease severity ratings in *Verticillium*-inoculated plants, indicating a potential for inducing systemic resistance, other aspects of the findings were inconclusive. Concentrations of terpenoid phytoalexins, indicative of a plant's defense response, were not consistently altered by *T. virens* treatment in *Verticillium*-inoculated plants. Moreover, the growth-promoting effects of *T. virens* were evident in the absence of *Vd*, suggesting a dual role of some *Trichoderma* strains [33]. These

aspects highlight the complexity of the interactions between *Trichoderma* strains, the target pathogen, and the surrounding environment. The variable outcomes across different studies, including ours, emphasize that the success of *Trichoderma*-based approaches against *Vd* is context-dependent. In contrast to *Trichoderma*, the literature suggests that *Clonostachys*-based treatments may not be the most suitable option for enhancing biological control of *Verticillium* wilt. While a few studies have noted the biocontrol impact of *Clonostachys* on *Verticillium* wilt, instances of successful *in planta* biocontrol reports remain scarce. Although certain *Clonostachys* strains have demonstrated the ability to decrease microsclerotia viability under soil conditions, the literature offers limited accounts of effective *in planta* biocontrol [34–37].

The Strepse formulation, comprising *Streptomyces* and *Pseudomonas* microorganisms, showed efficacy against *Vd* in controlled conditions (Figure 1a,b) but was less effective in the greenhouse (Figure 4a,b). Diverse *Streptomyces* species have been extensively investigated for their potential as biological control agents against *Verticillium*. Four strains isolated from diverse crop rhizospheres were assessed for their antagonistic impact on *Vd* in cotton, demonstrating biocontrol efficiencies ranging from 19% to 66% under greenhouse conditions, and slightly lower efficiencies between 14% and 51% in field conditions, dependent on application methods [38]. Co-inoculating *Arabidopsis thaliana* seeds with *Vd* and *Streptomyces lividans* 66 effectively suppressed the fungus in the soil, leading to reduced *Verticillium*-induced disease symptoms [39]. In greenhouse experiments with potato, tomato, and strawberry, various *Streptomyces* species decreased disease incidence and severity [40–43]. However, the biofungicide Mycostop R based on *S. griseovirides* K61 did not significantly protect tomato against *Vd* [44]. In addition to *Streptomyces*, *Pseudomonas* spp. show promise as biocontrol agents against various *Verticillium*-induced diseases in diverse crops, with a substantial volume of data supporting their potential efficacy [40,41,45–51]. Interestingly, while the results of our study indicated successful disease mitigation in controlled conditions for Strepse formulation, the efficacy was not consistently replicated in the greenhouse experiment which may be attributed to the complex and dynamic environmental factors present in that setting.

In the present study, Strepse did not exhibit effectiveness against *Fol*. This finding contrasts with previous research highlighting the efficacy of other *Streptomyces* isolates against this pathogen [39,47–50]. However, other studies have also reported limitations in the effectiveness of *Streptomyces* isolates [52,53]. These contrasting results suggest that the success of *Streptomyces* isolates against *Fol* may be influenced by various experimental conditions and contexts. Similarly, specific strains of *Pseudomonas* have been well acknowledged as effective biological control agents against numerous fungal pathogens, including *F. oxysporum* [54]. Extensive examples can be found in the literature demonstrating various *Pseudomonas* species safeguarding plants from *Fusarium* wilt, spanning tomato infections, and a variety of other plant hosts [55–58]. However, it is important to note that not all *Pseudomonas* strains exhibit effectiveness against *Fol* [59].

Our results of the pathogenicity experiments, both in the growth room and greenhouse, suggest promise for Clonotri as a biocontrol product for mitigating *Fol* infection. While Strepse did not reach statistical significance, it exhibits a trend toward reducing symptoms for *Vd*. The difference in efficacy of Strepse against *Vd* between controlled conditions and the greenhouse experiment can be attributed to several factors. Firstly, the controlled conditions in the initial experiment provided a more controlled and optimized environment for evaluating the biocontrol product. This controlled environment may have allowed for better establishment and performance of Strepse against *Vd*, resulting in significant protection observed. In contrast, the greenhouse experiment introduced more realistic and complex environmental conditions, including variations in temperature, humidity, and overall plant health. These factors can influence the interaction between the biocontrol product and the pathogen, potentially affecting the efficacy of Strepse. It is important to note that the lack of significant protection observed in the greenhouse experiment does not necessarily imply the complete ineffectiveness of Strepse against *Vd*. It could indicate that the

efficacy of Strepse is influenced by the specific conditions and dynamics of the greenhouse environment. Further investigations under various field conditions and with different cultivars are necessary to better understand the effectiveness of Strepse as a biocontrol agent against *Vd*.

This study also evaluated the potential synergistic effects of a combined application of Strepse and Clonotri. However, the combined treatment employed half the recommended dosage of each individual product. This reduced application rate was chosen to assess their effect in combination while avoiding potential overapplication effects. Nevertheless, this dilution might have resulted in a suboptimal concentration of the active microorganisms' spores, potentially limiting their efficacy against *Vd* and *Fol*. Further investigations are warranted to determine if a full recommended dose of the combined Strepse and Clonotri formulation would demonstrate improved biocontrol against *Vd* and *Fol*.

The present study also investigated the influence of pathogen concentration on biocontrol efficacy. The evaluation of biocontrol products against different pathogen concentrations in a controlled environment provides valuable insights into their efficacy across a range of disease pressures. Building upon the findings from the previous experiment, this phase aimed to understand how the biocontrol products Strepse and Clonotri perform when exposed to varying concentrations of *Vd* and *Fol*, respectively. In the case of Strepse against *Vd*, AUDPC analysis indicated significant protection at higher pathogen concentrations (10^6 and 10^5 conidia/mL), while its protective effect was not significant at a low concentration (10^4 conidia/mL) (Figure 2d). Similarly, Clonotri's efficacy against *Fol*, as revealed by AUDPC analysis, was prominent only at the highest pathogen concentration (10^6 conidia/mL), with no significant protection observed at lower concentrations (10^4 and 10^5 conidia/mL) (Figure 3d). The observed pattern of biocontrol effectiveness by Strepse and Clonotri may be related to the baseline disease severity at lower pathogen concentrations. Since plants without biocontrol products exhibited relatively low disease severity in these conditions, the impact of these treatments on disease reduction might not have been statistically significant. This suggests that the effectiveness of these biocontrol agents against their respective pathogens depends on the level of inoculum pressure. It has been extensively demonstrated that the severity of fungal wilt disease symptoms correlates with increasing inoculum concentrations. In a pertinent study conducted by Shaw et al. [60], strawberry genotypes previously classified as relatively susceptible, intermediate, and resistant to *Verticillium dahliae*-induced wilt were inoculated with varying conidial concentrations (10^4 , 10^5 , and 10^6 conidia/mL). Consistent with our own investigation, their results revealed a proportional increase in overall disease symptoms with rising inoculum concentrations. Specifically, the percentages of plants exhibiting stunting or mortality were 6.8%, 32.6%, and 44.9% for the three conidial concentrations, respectively. Furthermore, the study demonstrated that the categories of relative resistance were distinguishable at concentrations of 10^6 and 10^5 conidia/mL. However, no clear differentiation was observed at 10^4 conidia/mL, likely due to the relatively low disease severity at this specific concentration of the pathogen. It is worth mentioning that in our study, the assessment of plant height and fresh weight, essential indicators of plant health and growth, revealed that the application of biocontrol products did not have any discernible effect on these parameters, irrespective of whether the plants were part of the control group or were inoculated with a pathogen (Figures S3 and S4). Our results indicate that the protective potential of the biocontrol products is most pronounced when disease pressure is higher, demonstrating the interplay between biocontrol efficacy and baseline disease severity. Furthermore, the negligible impact on plant height and fresh weight emphasizes the lack of adverse effects on plant growth, affirming the potential of these products for sustainable disease management strategies.

In the context of plant growth and productivity, the application of biocontrol products did not significantly affect parameters like plant height, number of leaves, fruit number, and overall productivity under greenhouse conditions (Figures S5b,c and 5a,c). Interestingly, Clonotri-treated plants infected with *Fol* displayed no significant difference in mean fresh

weight compared to non-treated (-ve control) and Clonotri-treated control plants (without *Fol* infection) (Figure 5b). This suggests a potential protective effect of Clonotri treatment on this fruit yield parameter against *Fol* infection, which is consistent with the disease severity reduction observed in the pathogenicity experiment (Figure 4d).

The greenhouse experiment also included an evaluation of the influence of the biocontrol products on fruit quality attributes. Interestingly, the application of biocontrol products did not have a substantial impact on parameters such as titratable acidity, total phenolic and flavonoid contents, β -carotene and lycopene contents, and antioxidant activity (Figure S7). However, fruit firmness was increased in the first fruit set in *Fol*-infected plants compared to mock-inoculated plants, indicating that *Fol* infection might influence the fruit's ripening process (Figure 6). Similarly, the analysis of external fruit color revealed that *Fol* infection resulted in a lighter appearance of the fruit and a noticeable shift towards green and yellow coloration (Table 1). These effects are likely a result of *Fol*'s interference with the plant's vascular system, reduced water uptake, induced stress-related metabolic changes, and potential disruption of pigment synthesis pathways. Soluble solids content (SSC) was also influenced by pathogen infection, with *Vd*-infected plants displaying increased SSC, possibly indicating enhanced metabolic processes related to fruit ripening (Figure 7a,b). Verticillium wilt is known to impact the ripening process in infected plants, particularly in fruit-bearing crops like tomatoes. The induction of ripening in Verticillium-infected plants involves complex interactions between the pathogen and the host plant, including plant stimulation to produce ethylene, a key plant hormone involved in the initiation and acceleration of the ripening process [61].

5. Conclusions

This study investigated the efficacy of two commercially available biocontrol formulations, Strepse and Clonotri, against the fungal wilt pathogens *Verticillium dahliae* (*Vd*) and *Fusarium oxysporum* f. sp. *lycopersici* (*Fol*) in tomato cultivation. While Strepse showed a promising trend toward reducing *Vd* symptoms under controlled conditions, its efficacy in the greenhouse experiment was likely influenced by complex environmental factors. Conversely, Clonotri demonstrated significant disease mitigation against *Fol* in both controlled and greenhouse settings. The effectiveness of both biocontrol agents was dependent on pathogen concentration, with stronger protective effects observed at higher inoculum pressures. Importantly, the application of these products did not negatively impact plant growth or overall fruit quality, suggesting their potential for sustainable disease management strategies. These findings highlight the complexity of biocontrol efficacy, which is influenced by pathogen specificity, environmental factors, and plant health. Future research should investigate the performance of these biocontrol agents under diverse field conditions with different cultivars while also considering broader ecological interactions to fully harness their potential and optimize their use in practical tomato production. In conclusion, this study provides a valuable step forward in understanding the role of biocontrol products as a sustainable and environmentally friendly approach to disease management in tomato cultivation.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/agriculture14060882/s1>; Figure S1. Interaction plot showing disease severity on the y-axis and *Vd* concentration on the x-axis for both the treated (Strepse) and control groups; Figure S2. Interaction plot showing disease severity on the y-axis and *Fol* concentration on the x-axis for both the treated (Clonotri) and control groups; Figure S3: Growth characteristics in plants treated with biocontrol products and inoculated with varied *Vd* concentrations in growth room conditions; Figure S4: Growth characteristics in plants treated with biocontrol products and inoculated with varied *Fol* concentrations in growth room conditions; Figure S5: Growth characteristics in plants treated with biocontrol products and inoculated with the pathogens in greenhouse conditions; Figure S6: Fruit firmness of the second fruit set in plants inoculated with *V. dahliae* and *F. oxysporum* f. sp. *lycopersici* and treated with biocontrol products in the greenhouse experiment; Figure S7: Fruit quality attributes analyzed in two fruit sets of plants inoculated with *V. dahliae* and *F. oxysporum* f.

sp. *lycopersici* and treated with biocontrol products in the greenhouse experiment; Table S1: Mean values of color attributes (L^* , a^* , b^*) in tomato fruit of the second fruit set, as affected by the biocontrol products and the pathogens.

Author Contributions: Conceptualization, I.P. and M.-D.T.; methodology, I.P. and V.G.; validation, M.-D.T., I.P. and V.G.; formal analysis, I.P. and V.G.; investigation, G.D., S.P., M.-D.T., L.V. and I.P.; resources, I.P.; data curation, G.D., S.P., M.-D.T. and I.P.; writing—original draft preparation, I.P.; writing—review and editing, I.P., M.-D.T. and V.G.; visualization, I.P.; supervision, I.P. and V.G.; project administration, I.P. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Data Availability Statement: The data presented in this study are available on request from the corresponding author.

Acknowledgments: The authors thank Eleni Kiourtsidou and Apostolos Zavos for their expertise in the maintenance of the greenhouse facilities. The authors would like to acknowledge the support of T.P. Agrobiotech LTD for offering the commercial formulations Clonotri and Strepse. used for experiments.

Conflicts of Interest: The authors declare no conflicts of interest.

References

- Larena, I.; Sabuquillo, P.; Melgarejo, P.; De Cal, A. Biocontrol of Fusarium and Verticillium Wilt of Tomato by *Penicillium oxalicum* under Greenhouse and Field Conditions. *J. Phytopathol.* **2003**, *151*, 507–512. [[CrossRef](#)]
- De Jonge, R.; Van Esse, H.P.; Maruthachalam, K.; Bolton, M.D.; Santhanam, P.; Saber, M.K.; Zhang, Z.; Usami, T.; Lievens, B.; Subbarao, K.V.; et al. Tomato Immune Receptor Ve1 Recognizes Effector of Multiple Fungal Pathogens Uncovered by Genome and RNA Sequencing. *Proc. Natl. Acad. Sci. USA* **2012**, *109*, 5110–5115. [[CrossRef](#)] [[PubMed](#)]
- Di, X.; Gomila, J.; Takken, F.L.W. Involvement of Salicylic Acid, Ethylene and Jasmonic Acid Signalling Pathways in the Susceptibility of Tomato to *Fusarium oxysporum*. *Mol. Plant Pathol.* **2017**, *18*, 1024–1035. [[CrossRef](#)] [[PubMed](#)]
- Wang, M.; Sun, Y.; Sun, G.; Liu, X.; Zhai, L.; Shen, Q.; Guo, S. Water Balance Altered in Cucumber Plants Infected with *Fusarium oxysporum* f. Sp. *cucumerinum*. *Sci. Rep.* **2015**, *5*, 7722. [[CrossRef](#)] [[PubMed](#)]
- Fradin, E.F.; Thomma, B.P.H.J. Physiology and Molecular Aspects of Verticillium Wilt Diseases Caused by *V. dahliae* and *V. albo-atrum*. *Mol. Plant Pathol.* **2006**, *7*, 71–86. [[CrossRef](#)] [[PubMed](#)]
- Michielse, C.B.; Rep, M. Pathogen Profile Update: *Fusarium oxysporum*. *Mol. Plant Pathol.* **2009**, *10*, 311–324. [[CrossRef](#)]
- Antoniou, A.; Tsolakidou, M.-D.; Stringlis, I.A.; Pantelides, I.S. Rhizosphere Microbiome Recruited from a Suppressive Compost Improves Plant Fitness and Increases Protection against Vascular Wilt Pathogens of Tomato. *Front. Plant Sci.* **2017**, *8*, 2022. [[CrossRef](#)]
- Yadeta, K.A.; Thomma, B.P.H.J. The Xylem as Battleground for Plant Hosts and Vascular Wilt Pathogens. *Front. Plant Sci.* **2013**, *4*, 97. [[CrossRef](#)]
- Pascale, A.; Proietti, S.; Pantelides, I.S.; Stringlis, I.A. Modulation of the Root Microbiome by Plant Molecules: The Basis for Targeted Disease Suppression and Plant Growth Promotion. *Front. Plant Sci.* **2020**, *10*, 1741. [[CrossRef](#)]
- De Coninck, B.; Timmermans, P.; Vos, C.; Cammue, B.P.A.; Kazan, K. What Lies beneath: Belowground Defense Strategies in Plants. *Trends Plant Sci.* **2015**, *20*, 91–101. [[CrossRef](#)]
- Subbarao, K.V.; Kabir, Z.; Martin, F.N.; Koike, S.T. Management of Soilborne Diseases in Strawberry Using Vegetable Rotations. *Plant Dis.* **2007**, *91*, 964–972. [[CrossRef](#)] [[PubMed](#)]
- Song, Y.; Liu, L.; Wang, Y.; Valkenburg, D.J.; Zhang, X.; Zhu, L.; Thomma, B.P.H.J. Transfer of Tomato Immune Receptor Ve1 Confers Ave1-Dependent Verticillium Resistance in Tobacco and Cotton. *Plant Biotechnol. J.* **2018**, *16*, 638–648. [[CrossRef](#)] [[PubMed](#)]
- Saremi, H.; Okhovat, S.M.; Ashrafi, S.J. Fusarium Diseases as the Main Soil Borne Fungal Pathogen on Plants and Their Control Management with Soil Solarization in Iran. *Afr. J. Biotechnol.* **2011**, *10*, 18391–18398. [[CrossRef](#)]
- Thies, J.A. Grafting for Managing Vegetable Crop Pests. *Pest. Manag. Sci.* **2021**, *77*, 4825–4835. [[CrossRef](#)] [[PubMed](#)]
- Köhl, J.; Kolnaar, R.; Ravensberg, W.J. Mode of Action of Microbial Biological Control Agents against Plant Diseases: Relevance beyond Efficacy. *Front. Plant Sci.* **2019**, *10*, 845. [[CrossRef](#)] [[PubMed](#)]
- van Lenteren, J.C.; Bolckmans, K.; Köhl, J.; Ravensberg, W.J.; Urbaneja, A. Biological Control Using Invertebrates and Microorganisms: Plenty of New Opportunities. *BioControl* **2018**, *63*, 39–59. [[CrossRef](#)]
- Montesinos, E.; Bonaterra, A. Microbial Pesticides. In *Encyclopedia of Microbiology*; Schaechter, M., Ed.; Elsevier: Amsterdam, The Netherlands, 2009; pp. 110–120.
- Bonaterra, A.; Badosa, E.; Daranas, N.; Francés, J.; Roselló, G.; Montesinos, E. Bacteria as Biological Control Agents of Plant Diseases. *Microorganisms* **2022**, *10*, 1759. [[CrossRef](#)] [[PubMed](#)]

19. Pantelides, I.S.; Tjamos, S.E.; Paplomatas, E.J. Insights into the Role of Ethylene Perception in Tomato Resistance to Vascular Infection by *Verticillium dahliae*. *Plant Pathol.* **2010**, *59*, 130–138. [[CrossRef](#)]
20. Rep, M.; Meijer, M.; Houterman, P.M.; Van Der Does, H.C.; Cornelissen, B.J.C. *Fusarium oxysporum* Evades I-3-Mediated Resistance without Altering the Matching Avirulence Gene. *Mol. Plant-Microbe Interact.* **2005**, *18*, 15–23. [[CrossRef](#)]
21. Sinha, A.K.; Wood, R.K.S. Studies on the Nature of Resistance in Tomato Plants to *Verticillium albo-atrum*. *Ann. Appl. Biol.* **1968**, *62*, 319–327. [[CrossRef](#)]
22. Campbell, C.L.; Madden, L.V. *Introduction to Plant Disease Epidemiology*; John Wiley & Sons: New York, NY, USA, 1990; 728p.
23. González-Casado, S.; Martín-Belloso, O.; Elez-Martínez, P.; Soliva-Fortuny, R. Induced Accumulation of Individual Carotenoids and Quality Changes in Tomato Fruits Treated with Pulsed Electric Fields and Stored at Different Post-Treatments Temperatures. *Postharvest Biol. Technol.* **2018**, *146*, 117–123. [[CrossRef](#)]
24. Ochoa-Velasco, C.E.; Valadez-Blanco, R.; Salas-Coronado, R.; Sustaita-Rivera, F.; Hernández-Carlos, B.; García-Ortega, S.; Santos-Sánchez, N.F. Effect of Nitrogen Fertilization and Bacillus Licheniformis Biofertilizer Addition on the Antioxidants Compounds and Antioxidant Activity of Greenhouse Cultivated Tomato Fruits (*Solanum lycopersicum* L. Var. Sheva). *Sci. Hortic.* **2016**, *201*, 338–345. [[CrossRef](#)]
25. Goulas, V.; Georgiou, E. Utilization of Carob Fruit as Sources of Phenolic Compounds with Antioxidant Potential: Extraction Optimization and Application in Food Models. *Foods* **2020**, *9*, 20. [[CrossRef](#)]
26. Nagata, M.; Yamashita, I. Simple Method for Simultaneous Determination of Chlorophyll and Carotenoids in Tomato Fruit. *Jpn. Soc. Food Sci. Technol.* **1992**, *39*, 925–928. [[CrossRef](#)]
27. Larkin, R.P.; Fravel, D.R. Efficacy of Various Fungal and Bacterial Biocontrol Organisms for Control of Fusarium Wilt of Tomato. *Plant Dis.* **1998**, *82*, 1022–1028. [[CrossRef](#)] [[PubMed](#)]
28. Segarra, G.; Casanova, E.; Avilés, M.; Trillas, I. Trichoderma Asperellum Strain T34 Controls Fusarium Wilt Disease in Tomato Plants in Soilless Culture through Competition for Iron. *Microb. Ecol.* **2010**, *59*, 141–149. [[CrossRef](#)] [[PubMed](#)]
29. Cotxarrera, L.; Trillas-Gay, M.I.; Steinberg, C.; Alabouvette, C. Use of Sewage Sludge Compost and *Trichoderma asperellum* Isolates to Suppress Fusarium Wilt of Tomato. *Soil Biol. Biochem.* **2002**, *34*, 467–476. [[CrossRef](#)]
30. Sehimi, A.E.; Hewedy, O.A.; Altammar, K.A.; Alhumaidi, M.S.; Abd Elghaffar, R.Y. *Trichoderma asperellum* Empowers Tomato Plants and Suppresses *Fusarium oxysporum* through Priming Responses. *Front. Microbiol.* **2023**, *14*, 1140378. [[CrossRef](#)]
31. Høyer, A.K.; Jørgensen, H.J.L.; Amby, D.B.; Jensen, B. Endophytic Colonization of Tomato Plants by the Biological Control Agent *Clonostachys rosea*. In *Biological and Integrated Control of Plant Pathogens*; IOBC-WPRS Bulletin; Department of Plant and Environmental Sciences: Frederiksberg, Denmark, 2016; Volume 115, pp. 101–106.
32. Ordentlich, A.; Nachmias, A.; Chet, I. Integrated Control of Verticillium Dah/ / Ae in Potato by *Trichoderma harzlanum* and Captan. *Crop Prot.* **1990**, *9*, 363–366. [[CrossRef](#)]
33. Hanson, L.E. Plant Pathology & Nematology Reduction of Verticillium Wilt Symptoms in Cotton Following Seed Treatment with *Trichoderma virens*. *J. Cotton Sci.* **2000**, *4*, 224–231.
34. Varo, A.; Raya-Ortega, M.C.; Trapero, A. Selection and Evaluation of Micro-Organisms for Biocontrol of *Verticillium dahliae* in Olive. *J. Appl. Microbiol.* **2016**, *121*, 767–777. [[CrossRef](#)] [[PubMed](#)]
35. Alström, S. Root-Colonizing Fungi from Oilseed Rape and Their Inhibition of *Verticillium dahliae*. *J. Phytopathol.* **2000**, *148*, 417–423. [[CrossRef](#)]
36. Deketelaere, S.; Tyvaert, L.; França, S.C.; Hofte, M. Desirable Traits of a Good Biocontrol Agent against Verticillium Wilt. *Front. Microbiol.* **2017**, *8*, 1186. [[CrossRef](#)] [[PubMed](#)]
37. Keinath, A.P.; Fravel, D.R.; Papavizas, G.C. Potential of Gliocladium Roseum for Biocontrol of *Verticillium dahliae*. *Phytopathology* **1991**, *81*, 644–648. [[CrossRef](#)]
38. Xue, L.; Xue, Q.; Chen, Q.; Lin, C.; Shen, G.; Zhao, J. Isolation and Evaluation of Rhizosphere Actinomycetes with Potential Application for Biocontrol of Verticillium Wilt of Cotton. *Crop Prot.* **2013**, *43*, 231–240. [[CrossRef](#)]
39. Meschke, H.; Schrempf, H. Streptomyces Lividans Inhibits the Proliferation of the Fungus *Verticillium dahliae* on Seeds and Roots of *Arabidopsis thaliana*. *Microb. Biotechnol.* **2010**, *3*, 428–443. [[CrossRef](#)] [[PubMed](#)]
40. Berg, G.; Kurze, S.; Buchner, A.; Wellington, E.M.; Smalla, K.; Berg, G.; Kurze, S.; Wellington, E.M.; Buchner, A.; Smalla, K. Successful Strategy for the Selection of New Strawberry-Associated Rhizobacteria Antagonistic to Verticillium Wilt. *Can. J. Microbiol.* **2000**, *46*, 1128–1137. [[CrossRef](#)] [[PubMed](#)]
41. Berg, G.; Fritze, A.; Roskot, N.; Smalla, K. Evaluation of Potential Biocontrol Rhizobacteria from Different Host Plants of *Verticillium dahliae* Kleb. *J. Appl. Microbiol.* **2001**, *91*, 963–971. [[CrossRef](#)] [[PubMed](#)]
42. Entry, J.A.; Strasbaugh, C.A.; Sojka, R.E. Wood Chip-Polyacrylamide Medium for Biocontrol Bacteria Decreases *Verticillium dahliae* Infection on Potato. *Biocontrol Sci. Technol.* **2000**, *10*, 677–686. [[CrossRef](#)]
43. Cao, P.; Liu, C.; Sun, P.; Fu, X.; Wang, S.; Wu, F.; Wang, X. An Endophytic *Streptomyces* Sp. Strain DHV3-2 from Diseased Root as a Potential Biocontrol Agent against *Verticillium dahliae* and Growth Elicitor in Tomato (*Solanum lycopersicum*). *Antonie Leeuwenhoek Int. J. Gen. Mol. Microbiol.* **2016**, *109*, 1573–1582. [[CrossRef](#)]
44. Minuto, A.; Spadaro, D.; Garibaldi, A.; Gullino, M.L. Control of Soilborne Pathogens of Tomato Using a Commercial Formulation of *Streptomyces griseoviridis* and Solarization. *Crop Prot.* **2006**, *25*, 468–475. [[CrossRef](#)]
45. Leben, S.D.; Wadi, J.A.; Easton, G.D. Ecology and Epidemiology Effects of *Pseudomonas fluorescens* on Potato Plant Growth and Control of *Verticillium dahliae*. *Phytopathology* **1987**, *77*, 1592–1595. [[CrossRef](#)]

46. Mercado-Blanco, J.; Rodríguez-Jurado, D.; Hervás, A.; Jiménez-Díaz, R.M. Suppression of Verticillium Wilt in Olive Planting Stocks by Root-Associated Fluorescent *Pseudomonas* spp. *Biol. Control* **2004**, *30*, 474–486. [[CrossRef](#)]
47. Prieto, P.; Navarro-Raya, C.; Valverde-Corredor, A.; Amyotte, S.G.; Dobinson, K.F.; Mercado-Blanco, J. Colonization Process of Olive Tissues by *Verticillium dahliae* and Its in Planta Interaction with the Biocontrol Root Endophyte *Pseudomonas fluorescens* PICF7. *Microb. Biotechnol.* **2009**, *2*, 499–511. [[CrossRef](#)] [[PubMed](#)]
48. Malandraki, I.; Tjamos, S.E.; Pantelides, I.S.; Paplomatas, E.J. Thermal Inactivation of Compost Suppressiveness Implicates Possible Biological Factors in Disease Management. *Biol. Control* **2008**, *44*, 180–187. [[CrossRef](#)]
49. Uppal, A.K.; El Hadrami, A.; Adam, L.R.; Tenuta, M.; Daayf, F. Biological Control of Potato Verticillium Wilt under Controlled and Field Conditions Using Selected Bacterial Antagonists and Plant Extracts. *Biol. Control* **2008**, *44*, 90–100. [[CrossRef](#)]
50. Erdogan, O.; Benlioglu, K. Biological Control of Verticillium Wilt on Cotton by the Use of Fluorescent *Pseudomonas* Spp. under Field Conditions. *Biol. Control* **2010**, *53*, 39–45. [[CrossRef](#)]
51. Abuamsha, R.; Salman, M.; Ehlers, R.U. Differential Resistance of Oilseed Rape Cultivars (*Brassica napus* Ssp. *oleifera*) to *Verticillium longisporum* Infection Is Affected by Rhizosphere Colonisation with Antagonistic Bacteria, *Serratia plymuthica* and *Pseudomonas chlororaphis*. *BioControl* **2011**, *56*, 101–112. [[CrossRef](#)]
52. Tewfike, T.A. Biocontrol Activity of *Streptomyces Fulvoviolaceus* against *Fusarium oxysporum* and *Rhizoctonia solani* Pathogens of Tomato Plants. *Ann. Agric. Sci. Moshtohor* **2000**, *38*, 1503–1520.
53. Kanini, G.S.; Katsifas, E.A.; Savvides, A.L.; Karagouni, A.D. *Streptomyces Rochei* ACTA1551, an Indigenous Greek Isolate Studied as a Potential Biocontrol Agent against *Fusarium oxysporum* f. Sp. *lycopersici*. *Biomed. Res. Int.* **2013**, *2013*, 387230. [[CrossRef](#)]
54. Haas, D.; Défago, G. Biological Control of Soil-Borne Pathogens by Fluorescent *Pseudomonads*. *Nat. Rev. Microbiol.* **2005**, *3*, 307–319. [[CrossRef](#)] [[PubMed](#)]
55. Scher, F.M.; Baker, R. Mechanism of Biological Control in a Fusarium-Suppressive Soil. *Phytopathology* **1980**, *70*, 412–417. [[CrossRef](#)]
56. Toyoda, H.; Hashimoto, H.; Utsumi, R.; Kobayashi, H.; Ouchi, S. Detoxification of Fusaric Acid by a Fusaric Acid-Resistant Mutant of *Pseudomonas solanacearum* and Its Application to Biological Control of Fusarium Wilt of Tomato. *Phytopathology* **1988**, *78*, 1307–1311. [[CrossRef](#)]
57. van Peer, R.; Niemann, G.J.; Schippers, B. Induced Resistance and Phytoalexin Accumulation in Biological Control of Fusarium Wilt of Carnation by *Pseudomonas* Sp. Strain WCS417r. *Phytopathology* **1991**, *81*, 728–734. [[CrossRef](#)]
58. De Boer, M.; Bom, P.; Kindt, F.; Keurentjes, J.J.B.; Van Der Sluis, I.; Van Loon, L.C.; Bakker, P.A.H.M. Control of Fusarium Wilt of Radish by Combining *Pseudomonas putida* Strains That Have Different Disease-Suppressive Mechanisms. *Phytopathology* **2003**, *93*, 626–632. [[CrossRef](#)] [[PubMed](#)]
59. Lemanceau, P.; Alabouvette, C. Biological Control of Fusarium Diseases by Fluorescent *Pseudomonas* and Non-Pathogenic Fusarium. *Crop Prot.* **1991**, *10*, 279–286. [[CrossRef](#)]
60. Shaw, D.V.; Gubler, W.D.; Hansen, J. Field Resistance of California Strawberries to *Verticillium dahliae* at Three Conidial Inoculum Concentrations. *HortScience* **1997**, *32*, 711–713. [[CrossRef](#)]
61. Tsolakidou, M.-D.; Pantelides, I.; Tzima, A.K.; Kang, S.; Paplomatas, E.J.; Tsaltas, D. Disruption and Overexpression of the Gene Encoding ACC (1-Aminocyclopropane-1-Carboxylic Acid) Deaminase in Soil-Borne Fungal Pathogen *Verticillium dahliae* Revealed the Role of ACC as a Potential Regulator of Virulence and Plant Defense. *Mol. Plant-Microbe Interact.* **2019**, *32*, 639–653. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.